

LCGT F2F August 3–5, 2011

EM Counterpart Search for GW events

Nobuyuki Kawai

Department of Physics, Tokyo Tech

Outline



- Short GRB – usual suspect
 - Properties, short vs. long
 - In the context of standard paradigm
- Counterpart search strategy
 - Location: how good it should be
 - Follow up or coincidence?
- Facilities
- Robotic telescope

Why talk about GRB?

















- “Short GRB” is the prime candidate for EM counterpart of promised GW source (NS merger)
- Lessons from the history of GRB research
 - Cosmic source with poor localization
 - First detection (~1967, published in 1973)
 - First identification of counterpart (1997)

Brief review of GRBs (~1967–1997)

- Discovered in 60's, distance unknown for 30 years
- keV—GeV range (typ. hundreds of keV)
- Dominate the sky in X-ray and gamma-ray
- One GRB per day in the entire sky
- Isotropic distribution in the sky
- Not uniform in space: deficit of dim bursts
- Variety of light curves, short and long classes



x10³ Counts/s








Seconds (Rel. to Trigger)

Seconds (Rel. to Trigger)

Brief review of GRBs (1997~)

- Often associated with afterglow in X, optical, ...
- Associated with distant galaxies ($z \sim 1$ or larger)
- Fireball model favored
- Evidence for collimated outflow
- Long GRBs occur in star forming galaxies
- Some long GRBs associated with supernova
- Isotropic distribution in the sky
- Not uniform in space: deficit of dim bursts




Fireball scenario of GRBs

Paczynski, Meszaros, Rees, Sari, Piran, ...



GRB Alert Network



GRB



- Classification: long vs. short
- Identification of GRBs
 - Early days –no success
 - Modulation collimator, aspect ratio: degree, and slow
 - Interplanetary network: arcmin, months
 - BACODINE: 5 degrees, fast (20 seconds),
 - Afterglow detection
- First in X-ray
 - brightest source in the sky: easy to find
 - Imaging available: arcmin (coded mask) ~ arcsec (focusing mirror)
- Following up in optical
 - Usually faint, >20 mag at 1 day
 - » No catalog (maybe SDSS)
 - about Half obscured in optical (IR?)
 - could be bright at very early phase (<1 min of GRB)

Example of follow-up observations

- GRB 970228
 - GRBM → WFC –(human) → MECS (X-ray telescope) – fading source
 - → Optical telescope – fading source
- GRB 990123
 - BATSE –(BACODINE)- ROTSE
 - GRBM/WFC –(human) → MECS (XRT) → ROTSE inspection
- GRB 030329
 - HETE-2 → Tokyo Tech 30cm
- GRB050509B
 - Swift BAT-XRT → Subaru etc.
- GRB050709
 - HETE-2 – Chandra – Hubble
- GRB070201
 - IPN – optical/LIGO
- GRB110721A
 - Fermi/LAT $D=1.5$ deg, 11 hours after the burst
 - Swift/XRT($D=0.4$ deg) incomplete tiling
 - Candidate X-ray/optical source found 2 days later
 - Turned out to be an early-type galaxy


GRB970228: first afterglow

- Coincident detection in gamma and X rays
- X-ray position analyzed on ground
- Satellite repointed in 8 hours
- X-ray afterglow imaged by focusing instrument



8 hours


3 days




Costa et al. 1997

GRB 970228: Discovery of optical transient

- Optical observation conducted based on WFC position (\sim a few arcmin)
- Fading source detected
- Later Hubble observation revealed underlying extended source, probably the host of the GRB




van Paradijs et al. 1997





GRB 990123

- ROTSE responded to BATSE ~ 5 degree localization
- 9 mag OT found in the BeppoSAX-WFC location (\sim a few arcmin)



GRB 030329 – Localization by HETE-2




Trigger 11:37:14
 Alert 11:38:41
 SXC 12:50:24 (+73 min)

OT 12:52:09 (+75 min)
 Torii GCN #1986
 Peterson
 GCN#1985




- WXM 1-D localization
- SXC 1 arcmin position in ~1hour with ground analysis
- 12-13 mag optical counterpart discovered immediately

GRB 030329



Observation with Tokyo Tech 30cm



Light curve of the optical afterglow of GRB 030329




GRB 050509B – first short GRB with arcmin localization



Gehrels et al. 2005


- Swift BAT → X-ray afterglow by Swift XRT
- Association with an elliptical galaxy at $z=0.225$: likely, but not conclusive






Subaru (Kosugi, et al. 2005)

GRB050709

short GRB localized by HETE-2




<http://space.mit.edu/HETE/Bursts/GRB050709/>




GRB 050709:

18

localization of X-ray and optical afterglow



Chandra X-ray Observatory



Hubble Space Telescope

- HETE WXM+SXC ~ 1 arcmin
- Chandra ~ 0.5 arcsec
- Hubble localized it to a dwarf starforming galaxy



$z=0.160$

Dwarf irregular galaxy

$\text{SFR} = 0.2 M_{\text{sun}}/\text{yr}$

GRB070201

– short GRB from Andromeda galaxy? (780 kpc)?





- LIGO measurements indicate that this could not have been a binary merger in M31 (Abbott et al. 2008)

$$E_{\gamma} = 1.5 \times 10^{45} \text{ erg}$$

GRB 110721A – GeV localization

- Fermi/LAT D=1.5 deg, 11 hours after the burst (large error and slow)
- Swift/XRT(D=0.4 deg) incomplete tiling
- Candidate X-ray/optical source found 2 days later
- Turned out to be an early-type galaxy



Lessons

- Pre-afterglow Era (~1997)
 - IPN: arcmin position is useless if too late
 - BATSE: prompt degrees position is useful with more constraining information
- Afterglow era (1997~)
 - Quick location is the key
 - Optical astronomers do not look at error circles larger than their field of view
 - Swift XRT: even arcmin position does not give conclusive identification
 - Optical/Near infrared imaging/spectroscopy required for unambiguous identification


Recommendations

- Localization error – the smaller the better
- Notification delay - the shorter the better

“Short hard” GRB


No afterglow detected
(until 2005)

- Different origin
 - supernovae?
 - neutron star merger?
 - magnetar flare?



Prompt emission of long and short GRBs

24



Delay in HE onset: 0.1-0.2 s


Delay in HE onset: ~4-5 s

HE delayed onset can be seen from almost all LAT GRBs


Afterglow of short and long GRBs

Short GRB050724

Long GRBs



Barthelmy et al. 2005






Nousek et al. 2006


X-ray afterglows are similar to those of long GRBs

Suspect –Merging neutron star binary


J.H. Taylor and R.H Hulse
1993 Nobel Prize in Physics



Known neutron star binaries




Redshift, isotropic radiated energy, and X-ray afterglow luminosity of GRBs (arrows: short GRBs)



Host galaxy and redshift of short GRBs

Optical Afterglows X-ray Afterglows


Berger et al. 2007; Berger 2009



E: elliptical galaxies (old stars only)

F: Star forming galaxies

Confirmed hosts – E:SF = 2:1




Half of short GRB at
 $z > 0.7$


$\Rightarrow \langle \text{age} \rangle \leq 7 \text{ Gyr}$

Star formation rate and metallicity of short GRB hosts

Berger 2009




Short GRB hosts have lower specific star formation rates than long GRB hosts; they trace the general galaxy population




Short GRB hosts have higher metallicities than long GRB hosts; they trace the general galaxy population

Sites of short GRBs in host galaxy



Short GRBs have larger offset from the center of their host galaxies




Short GRBs trace the light distribution of their host galaxies

Short GRB

- Occur in every type of galaxies
 - Traces star density
- Fermi LAT observation of GRB 090510:
 - high Lorentz factor – probably strong collimation
- Neutron star merger scenario favored, but not conclusive
 - Final word only given by GW detection
- If collimated (likely), there should be many off-axis events.
 - Orphan afterglow
 - GW events relatively close-by → bright orphan afterglow
 - follow up to GW trigger (large error box)
 - → continuous monitor of large sky region for short transients


Detection rate of neutron star merger by GW telescope

- LIGO (range ~ 20 Mpc)
→ on per 10–630 year (95%CL)
- LCGT (range ~ 200 Mpc:
 $z=0.015$)
→ 2–100 per year
- Short GRB detections <100 /year
 - Most DNS mergers are not detected as short GRB




Optical afterglow of GRBs

Short GRBs **with** measured **redshifts**




Short GRBs **without** measured **redshifts**





Optical afterglow of SGRB at $z=1$ and at 100 Mpc

35



Kann et al. 2011


Signature of jet break



Beamed emission. A relativistic jet with a Lorentz factor γ and an opening angle θ moves forward until its Lorentz factor $\gamma = \theta^{-1}$. Then it expands sideways rapidly, resulting in a "jet break" in the light curve. A schematic light curve is depicted at the top right.

Orphan afterglow for off-axis GRBs

Granot et al. 2011




Prediction for jetted GRBs

Many more orphan afterglows,
but fainter by order(s) of magnitude

Is neutron star merger detectable as a short GRB?


- GW telescope localization ≈ 10 deg
 - Extremely difficult to find optical counterpart
 - Error circle too large for “normal” telescopes
 - however, $z < 0.015 \rightarrow$ much closer than usual GRBs
 - \rightarrow need different strategy for counterpart search
- (1) Cover the large error circle (≥ 10 deg)
- (2) Continuous monitor of large sky
 - Find temporary coincidence with GW event
 - Search for orphan afterglow

Counterpart search strategy





Steps of localization and identification of GW events


Smaller error circle by EM



Detailed observation



Identification to astrophysical event



Akeno MITSuME 50 cm Telescope



50 cm autonomous robotic telescope at Akeno



8/4/11